

# The Handy Bt Trait Table

## for U.S. Corn Production

Updated  
August 2017

Posted at [www.msuent.com](http://www.msuent.com)

**VOID**

For questions, complaints, or corrections: Chris DiFonzo, Michigan State University, difonzo@msu.edu  
Contributors: Pat Porter, Texas A&M University & Kelley Tilmon, The Ohio State University

Most corn hybrids planted in the U.S. contain one or more transgenic traits for weed or insect management. These traits can increase flexibility and profitability for producers, but sometimes cause confusion about their spectrum of control or refuge requirements. The Handy Bt Trait Table provides a helpful list of trait names (below) and details of trait packages (next page) to make it easier to read company seed guides, sales materials, and bag tags. Note that there are two versions of the table (north/Midwest vs. south/cotton belt) which differ only in refuge percentages.

### Important clarifications or changes to the Trait Table for 2017

- ✓ An insect is listed in the CONTROL SPECTRUM column if seed providers claim protection or efficacy for a given Bt package; insect species which are 'suppressed' are no longer listed. Actual field-level performance of hybrids on lepidopteran and rootworm larvae may differ if there are local or regional insect populations which are less susceptible or resistant to Bt proteins.
- ✓ To address local or regional performance issues, a new column ('May be ineffective on') was added to highlight insect x Bt toxins with documented field-failures, confirmed resistance, or cross-resistance. An insect is listed in this column only if ALL of the Bt proteins which should control it in a product are 'ineffective' somewhere in the US or Canada. Ineffective ratings are based on published lab assays &/or field research from field corn, sweet corn, and cotton. University extension specialists or local educators can assist in determining if you are in an area where reduced effectiveness was reported. On a broader scale, this column is intended to alert growers and consultants to potential management problems, influence seed selection, and encourage field scouting.
- ✓ The refuge column was simplified to include only the % and an indication if the refuge is in the bag.


### Field corn 'events' (transformations of one or more genes) and their Trade Names

Trade name for trait	Event	Protein(s) expressed	Insect Target or Herbicide Activity
Agrisure CB/LL	Bt11	Cry1Ab + PAT	corn borer + <i>glufosinate tolerance</i>
Agrisure Duracade	5307	eCry3.1Ab	rootworm
Agrisure GT	GA21	EPSPS	<i>glyphosate tolerance</i>
Agrisure RW	MIR604	mCry3A	rootworm
Agrisure Viptera	MIR162	Vip3A	broad lep control (but not corn borer)
Herculex 1 or CB	TC1507	Cry1Fa2 + PAT	corn borer + <i>glufosinate tolerance</i>
Herculex RW	DAS-59122-7	Cry34Ab1/Cry35Ab1 + PAT	rootworm + <i>glufosinate tolerance</i>
(Not available – in Qrome)	4114	Cry1F + Cry34Ab1/Cry35Ab1 + PAT	corn borer+ rootworm+ <i>glufosinate tol.</i>
Roundup Ready 2	NK603	EPSPS	<i>glyphosate tolerance</i>
Yieldgard Corn Borer	MON810	Cry1Ab	corn borer
Yieldgard Rootworm	MON863	Cry3Bb1	rootworm
Yieldgard VT Pro	MON89034	Cry1A.105 + Cry2Ab2	broader lep control
Yieldgard VT Rootworm RR	MON88017	Cry3Bb1 + EPSPS	rootworm + <i>glyphosate tolerance</i>

### Abbreviations used in the Trait Table

#### Herbicide activity

- GT *glyphosate tolerant*
- LL Liberty Link - *glufosinate-tolerant*
- RR2 Roundup Ready 2, *glyphosate-tolerant*



#### Insect targets

- |                         |                              |
|-------------------------|------------------------------|
| BCW black cutworm       | SB stalk borer               |
| CEW corn earworm        | SCB sugarcane borer          |
| ECB European corn borer | SWCB southwestern corn borer |
| FAW fall armyworm       | TAW true armyworm            |
| RW corn rootworm        | WBC western bean cutworm     |



**Bt corn trait packages with their Bt proteins, spectrum of control, & % refuge**

Updated August 2017

TRAIT FAMILY	Specific Product	VOID Bt protein(s)	CONTROL SPECTRUM Marketed for control of: above-ground-----in soil		May be locally or regionally ineffective on:	Herbicide tolerance	Refuge %	
<b>AGRISURE</b>								
Agrisure 3010, 3010A		Cry1Ab	ECB	SCB	SWCB	---	GT LL 20%	
Agrisure 3000GT, 3011A		Cry1Ab mCry3A	ECB	SCB	SWCB	RW	RW GT LL 20%	
Agrisure Viptera 3110		Cry1Ab Vip3A	BCW	CEW	ECB	FAW	---	GT LL 20%
Agrisure Viptera 3111		Cry1Ab Vip3A mCry3A	BCW	CEW	ECB	FAW	RW	RW GT LL 20%
Agrisure 3120 E-Z Refuge		Cry1Ab Cry1F	BCW	ECB	FAW	---	FAW, WBC	REFER TO BAG TAG
Agrisure 3122 E-Z Refuge		Cry1Ab Cry1F mCry3A Cry34/35Ab1	BCW	ECB	FAW	---	FAW, WBC	for specific letter code:
Agrisure Viptera 3220 E-Z Refuge		Cry1Ab Cry1F Vip3A	BCW	CEW	ECB	FAW	---	EZ0 hybrids = GT only
Agrisure Duracade 5122 E-Z Refuge		Cry1Ab Cry1F mCry3A eCry3.1Ab	BCW	ECB	FAW	---	FAW, WBC	5% in bag
Agrisure Duracade 5222 E-Z Refuge		Cry1Ab Cry1F Vip3A mCry3A eCry3.1Ab	BCW	CEW	ECB	FAW	RW	EZ1 hybrids = GT LL
<b>HERCULEX</b>								
Herculex 1 (HX1)		Cry1F	BCW	ECB	FAW	---	FAW, SWCB, WBC	LL 20%
Herculex RW (HXRW)		Cry34/35Ab1	---	---	---	RW	RW	RR2 20%
Herculex XTRA (HXX)		Cry1F Cry34/35Ab1	BCW	ECB	FAW	---	FAW, SWCB, WBC	(most) 20%
<b>OPTIMUM</b>								
Intrasect (YHR)		Cry1Ab Cry1F	BCW	ECB	FAW	---	FAW, WBC	LL RR2 5%
AcreMax (AM)		Cry1Ab Cry1F	BCW	ECB	FAW	---	FAW, WBC	LL RR2 5% in bag
Leptra (VYHR) <sup>a</sup> AcreMax Leptra (AML) <sup>b</sup>		Cry1Ab Cry1F Vip3A	BCW	CEW	ECB	FAW	---	LL RR2 <sup>a</sup> 5% ½ mile <sup>b</sup> 5% in bag
AcreMax RW (AMRW)		Cry34/35Ab1	---	---	---	RW	RW	LL RR2 10% in bag
AcreMax1 (AM1)		Cry1F Cry34/35Ab1	BCW	ECB	FAW	---	FAW, SWCB, WBC	LL RR2 10% in bag 20% ECB
TRIssect (CHR)		Cry1F mCry3A	BCW	ECB	FAW	---	FAW, SWCB, WBC	LL RR2 20%
Intrasect TRIssect (CYHR) <sup>a</sup> AcreMax TRIssect (AMT) <sup>b</sup>		Cry1Ab Cry1F mCry3A	BCW	ECB	FAW	---	FAW, WBC	LL RR2 <sup>a</sup> 20% <sup>b</sup> 10% in bag
Intrasect Xtra (YXR) <sup>a</sup> AcreMax Xtra (AMX) <sup>b</sup>		Cry1Ab Cry1F Cry34/35Ab1	BCW	ECB	FAW	---	FAW, WBC	LL RR2 <sup>a</sup> 20% <sup>b</sup> 10% in bag
Intrasect Xtreme (CYXR) <sup>a</sup> AcreMax Xtreme (AMXT) <sup>b</sup>		Cry1Ab Cry1F mCry3A Cry34/35Ab1	BCW	ECB	FAW	---	FAW, WBC	LL RR2 <sup>a</sup> 5% <sup>b</sup> 5% in bag
<b>YIELDGARD or GENUITY</b>								
YieldGard CB (YGCB)		Cry1Ab	ECB	SCB	SWCB	---	SCB	RR2 20%
YieldGard VT Rootworm		Cry3Bb1	---	---	---	RW	RW	RR2 20%
YieldGard VT Triple		Cry1Ab Cry3Bb1	ECB	SCB	SWCB	---	SCB RW	RR2 20%
Genuity VT Double PRO <sup>a</sup> or RIB complete <sup>b</sup>		Cry1A.105 Cry2Ab2	CEW	ECB	FAW	---	CEW	RR2 <sup>a</sup> 5% <sup>b</sup> 5% in bag
Genuity VT Triple PRO <sup>a</sup> or RIB complete <sup>b</sup>		Cry1A.105 Cry2Ab2 Cry3Bb1	CEW	ECB	FAW	---	CEW RW	RR2 <sup>a</sup> 20% <sup>b</sup> 10% in bag
Genuity SmartStax <sup>a</sup> or RIB Complete <sup>b</sup>		Cry1A.105 Cry2Ab2 Cry1F Cry3Bb1 Cry34/35Ab1	BCW	CEW	ECB	FAW	RW	LL RR2 <sup>a</sup> 5% <sup>b</sup> 5% in bag
<b>OTHER</b>								
Powercore <sup>a</sup> Powercore Refuge Adv. <sup>b</sup>		Cry1A.105 Cry2Ab2 Cry1F	BCW	CEW	ECB	FAW	---	CEW, WBC
Smartstax <sup>a</sup> Smartstax Refuge Adv. <sup>b</sup>		Cry1A.105 Cry2Ab2 Cry1F Cry3Bb1 Cry34/35Ab1	BCW	CEW	ECB	FAW	RW	CEW, WBC RW
Qrome		Cry1Ab Cry1F mCry3A Cry34/35Ab1	BCW	ECB	FAW	---	FAW, WBC	LL RR2 5% in bag